

# Literaturhinweise

## Intrauterine fetale Transfusionen Intrauterine Transfusion von Erythrozyten

- 1) Liley AW.  
*Intrauterine transfusion of foetus in haemolytic disease.*  
*Br Med J* 1963; **2**:1107.
- 2) Rodeck CH, Kemp JR, Holman CA, et al.  
*Direct intravascular fetal blood transfusion by fetoscopy in severe Rhesus isoimmunisation.*  
*Lancet* 1981; **1**:625.
- 3) Nicolaides KH, Soothill PW, Clewell WH, et al.  
*Fetal haemoglobin measurement in the assessment of red cell isoimmunisation.*  
*Lancet* 1988; **1**:1073.
- 4) Mari G, Deter RL, Carpenter RL, et al.  
*Noninvasive diagnosis by Doppler ultrasonography of fetal anemia due to maternal red-cell alloimmunization. Collaborative Group for Doppler Assessment of the Blood Velocity in Anemic Fetuses.*  
*N Engl J Med* 2000; **342**:9.
- 5) von Kaisenberg CS, Jonat W.  
*Fetal parvovirus B19 infection.*  
*Ultrasound Obstet Gynecol* 2001; **18**:280.
- 6) Carr S, Rubin L, Dixon D, et al.  
*Intrauterine therapy for homozygous alpha-thalassemia.*  
*Obstet Gynecol* 1995; **85**:876.
- 7) Ng PC, Fok TF, Lee CH, et al.  
*Is homozygous alpha-thalassaemia a lethal condition in the 1990s?*  
*Acta Paediatr* 1998; **87**:1197.
- 8) Hayward A, Ambruso D, Battaglia F, et al.  
*Microchimerism and tolerance following intrauterine transplantation and transfusion for alpha-thalassemia-1.*  
*Fetal Diagn Ther* 1998; **13**:8.
- 9) Wang C, Ryan G.  
*Transfusion medicine illustrated: Intrauterine transfusion for homozygous alpha(0) thalassemia reverses hydrops fetalis.*  
*Transfusion* 2009; **49**:1043.
- 10) Remacha AF, Badell I, Pujol-Moix N, et al.  
*Hydrops fetalis-associated congenital dyserythropoietic anemia treated with intrauterine transfusions and bone marrow transplantation.*  
*Blood* 2002; **100**:356.
- 11) Ogburn PL Jr, Ramin KD, Danilenko-Dixon D, et al.  
*In utero erythrocyte transfusion for fetal xerocytosis associated with severe anemia and non-immune hydrops fetalis.*  
*Am J Obstet Gynecol* 2001; **185**:238.
- 12) Quarello E, Stirnemann J, Nassar M, et al.  
*Outcome of anaemic monochorionic single survivors following early intrauterine rescue transfusion in cases of feto-fetal transfusion syndrome.*  
*BJOG* 2008; **115**:595.
- 13) Quarello E, Stirnemann J, Nassar M, et al.  
*Outcome of anaemic monochorionic single survivors following early intrauterine rescue transfusion in cases of feto-fetal transfusion syndrome.* *BJOG* 2008; **115**:595.

- 14) Viëtor HE, Kanhai HH, Brand A.  
*Induction of additional red cell alloantibodies after intrauterine transfusions.*  
*Transfusion 1994; 34:970.*
- 15) Schonewille H, Klumper FJ, van de Watering LM, et al.  
*High additional maternal red cell alloimmunization after Rhesus- and K-matched intrauterine intravascular transfusions for hemolytic disease of the fetus.*  
*Am J Obstet Gynecol 2007; 196:143.e1.*
- 16) Watson WJ, Wax JR, Miller RC, Brost BC.  
*Prevalence of new maternal alloantibodies after intrauterine transfusion for severe Rhesus disease.*  
*Am J Perinatol 2006; 23:189.*
- 17) el-Azeem SA, Samuels P, Rose RL, et al.  
*The effect of the source of transfused blood on the rate of consumption of transfused red blood cells in pregnancies affected by red blood cell alloimmunization.*  
*Am J Obstet Gynecol 1997; 177:753.*
- 18) Dodd JM, Windrim RC, van Kamp IL.  
*Techniques of intrauterine fetal transfusion for women with red-cell isoimmunisation for improving health outcomes.*  
*Cochrane Database Syst Rev 2012; 9:CD007096.*
- 19) Harman CR, Bowman JM, Manning FA, Menticoglu SM.  
*Intrauterine transfusion--intraperitoneal versus intravascular approach: a case-control comparison.*  
*Am J Obstet Gynecol 1990; 162:1053.*
- 20) Lewis M, Bowman JM, Pollock J, Lowen B.  
*Absorption of red cells from the peritoneal cavity of an hydropic twin.*  
*Transfusion 1973; 13:37.*
- 21) Creasman WT, Duggan ER, Lund CJ.  
*Absorption of transfused chromium-labeled erythrocytes from the fetal peritoneal cavity in hydrops fetalis.*  
*Am J Obstet Gynecol 1966; 94:586.*
- 22) Taylor WW, Scott DE, Pritchard JA.  
*Fate of compatible adult erythrocytes in the fetal peritoneal cavity.*  
*Obstet Gynecol 1966; 28:175.*
- 23) Westgren M, Selbing A, Stangenberg M.  
*Fetal intracardiac transfusions in patients with severe rhesus isoimmunisation.*  
*Br Med J (Clin Res Ed) 1988; 296:885.*
- 24) Weiner CP, Wenstrom KD, Sipes SL, Williamson RA.  
*Risk factors for cordocentesis and fetal intravascular transfusion.*  
*Am J Obstet Gynecol 1991; 165:1020.*
- 25) Van Kamp IL, Klumper FJ, Oepkes D, et al.  
*Complications of intrauterine intravascular transfusion for fetal anemia due to maternal red-cell alloimmunization.*  
*Am J Obstet Gynecol 2005; 192:171.*
- 26) Nicolini U, Santolaya J, Ojo OE, et al.  
*The fetal intrahepatic umbilical vein as an alternative to cord needling for prenatal diagnosis and therapy.*  
*Prenat Diagn 1988; 8:665.*
- 27) Giannakopoulos X, Sepulveda W, Kourtis P, et al.  
*Fetal plasma cortisol and beta-endorphin response to intrauterine needling.*  
*Lancet 1994; 344:77.*

# Literaturhinweise

- 28)** Moise KJ Jr, Carpenter RJ Jr, Kirshon B, et al.  
*Comparison of four types of intrauterine transfusion: effect on fetal hematocrit.*  
*Fetal Ther 1989; 4:126.*
- 29)** Nicolini U, Kochenour NK, Greco P, et al.  
*When to perform the next intra-uterine transfusion in patients with Rh allo-immunization: combined intravascular and intraperitoneal transfusion allows longer intervals.*  
*Fetal Ther 1989; 4:14.*
- 30)** Fox C, Martin W, Somerset DA, et al.  
*Early intraperitoneal transfusion and adjuvant maternal immunoglobulin therapy in the treatment of severe red cell alloimmunization prior to fetal intravascular transfusion.*  
*Fetal Diagn Ther 2008; 23:159.*
- 31)** Forestier F, Daffos F, Catherine N, et al.  
*Developmental hematopoiesis in normal human fetal blood.*  
*Blood 1991; 77:2360.*
- 32)** Dildy GA 3rd, Smith LG Jr, Moise KJ Jr, et al.  
*Porencephalic cyst: a complication of fetal intravascular transfusion.*  
*Am J Obstet Gynecol 1991; 165:76.*
- 33)** Drew JH, Guarani RL, Cichello M, Hobbs JB.  
*Neonatal whole blood hyperviscosity: the important factor influencing later neurologic function is the viscosity and not the polycythemia.*  
*Clin Hemorheol Microcirc 1997; 17:67.*
- 34)** Mandelbrot L, Daffos F, Forestier F, et al.  
*Assessment of fetal blood volume for computer-assisted management of in utero transfusion.*  
*Fetal Ther 1988; 3:60.*
- 35)** Bowman JM.  
*The management of Rh-Isoimmunization.*  
*Obstet Gynecol 1978; 52:1.*
- 36)** Canlorbe G, Macé G, Cortey A, et al.  
*Management of very early fetal anemia resulting from red-cell alloimmunization before 20 weeks of gestation.*  
*Obstet Gynecol 2011; 118:1323.*
- 37)** Lobato G, Soncini CS.  
*Fetal hydrops and other variables associated with the fetal hematocrit decrease after the first intrauterine transfusion for red cell alloimmunization.*  
*Fetal Diagn Ther 2008; 24:349.*
- 38)** Mari G, Detti L, Oz U, et al.  
*Accurate prediction of fetal hemoglobin by Doppler ultrasonography.*  
*Obstet Gynecol 2002; 99:589.*
- 39)** Scheier M, Hernandez-Andrade E, Fonseca EB, Nicolaides KH.  
*Prediction of severe fetal anemia in red blood cell alloimmunization after previous intrauterine transfusions.*  
*Am J Obstet Gynecol 2006; 195:1550.*
- 40)** Mari, G, Zimmerman, R, Segata, M.  
*Am J Obstet Gynecol 2005; 191:S149.*

- 41) Trevett TN Jr, Dorman K, Lamvu G, Moïse KJ Jr.  
*Antenatal maternal administration of phenobarbital for the prevention of exchange transfusion in neonates with hemolytic disease of the fetus and newborn.*  
*Am J Obstet Gynecol 2005; 192:478.*
- 42) Watts DH, Luthy DA, Benedetti TJ, et al.  
*Intraperitoneal fetal transfusion under direct ultrasound guidance.*  
*Obstet Gynecol 1988; 71:84.*
- 43) Van Kamp IL, Klumper FJ, Oepkes D, et al.  
*Complications of intrauterine intravascular transfusion for fetal anemia due to maternal red-cell alloimmunization.*  
*Am J Obstet Gynecol 2005; 192:171.*
- 44) Lindenburg IT, Smits-Wintjens VE, van Klink JM, et al.  
*Long-term neurodevelopmental outcome after intrauterine transfusion for hemolytic disease of the fetus/newborn: the LOTUS study.*  
*Am J Obstet Gynecol 2012; 206:141.e1.*