

Molekulare Diagnostik in der Immunhämatologie - Wenn die Serologie nicht mehr ausreicht

1. Wagner FF, Frohmajer A, Flegel WA. RHD positive haplotypes in D negative Europeans. *BMC Genet* 2001;2:10.
2. Singleton BK, Green CA, Avent ND, Martin PG, Smart E, Daka A, Narter-Oлага EG, Hawthorne LM, Daniels G. The presence of an RHD pseudogene containing a 37 base pair duplication and a nonsense mutation in africans with the Rh D-negative blood group phenotype. *Blood* 2000;95:12-8.
3. Wagner FF, Flegel WA. RHD gene deletion occurred in the Rhesus box. *Blood* 2000;95:3662-8.
4. Curtis BR and McFarland JG. Human platelet antigens - 2013. *Vox Sang.* 2014;106:93-102.
5. Bux J. The human neutrophil antigen system: structure, function, and disease implications. *Hematology Education: the education program for the annual congress of the European Hematology Association.* 2015;9:401-7
6. Wagner FF, Gassner C, Müller TH, Schönitzer D, Schunter F, Flegel WA. Molecular basis of weak D phenotypes. *Blood.* 1999 Jan 1;93(1):385-93
7. Sandler SG, Flegel WA, Westhoff CM, Denomme GA, Delaney M, Keller MA, Johnson ST, Katz L, Queenan JT, Vassallo RR, Simon CD; College of American Pathologists Transfusion Medicine Resource Committee Work Group. It's time to phase in RHD genotyping for patients with a serologic weak D phenotype. *College of American Pathologists Transfusion Medicine Resource Committee Work Group. Transfusion.* 2015 Mar;55(3):680-9.
8. Daniels G. Variants of RhD--current testing and clinical consequences. *Br J Haematol.* 2013 May;161(4):461-70
9. Kacker S, Vassallo R, Keller MA, Westhoff CM, Frick KD, Sandler SG, Tobian AA. Financial implications of RHD genotyping of pregnant women with a serologic weak D phenotype. *Transfusion.* 2015 Mar 21. doi: 10.1111/trf.13074
10. Lo YMD, Corbetta N, Chamberlain PF, Rai V, Sargent IL, Redman CWG, Wainscoat JS. Presence of fetal DNA in maternal plasma and serum. *The Lancet* 1997;350: 485-487.
11. Chan KCA, Ding C, Gerovassili A, Yeung SW, Chiu RWK, Leung TN, Lau TK, Chim SSC, Chung GTY, Nicolaides KH, Lo YMD. Hypermethylated RASSF1A in Maternal Plasma: A Universal Fetal DNA Marker that Improves the Reliability of Noninvasive Prenatal Diagnosis. *Clin Chem* 2006;52: 2211-2218.
12. Doescher A, Petershofen EK, Wagner FF, Schunter M, Muller TH. Evaluation of single-nucleotide polymorphisms as internal controls in prenatal diagnosis of fetal blood groups. *Transfusion* 2013;53: 353-62.
13. Page-Christiaens GC, Bossers B, Van Der Schoot CE, De Haas M. Use of Bi-Allelic Insertion/Deletion Polymorphisms as a Positive Control for Fetal Genotyping in Maternal Blood. *Annals of the New York Academy of Sciences* 2006;1075: 123-129.
14. Wagner FF. RHD PCR of D-Negative Blood Donors. *Transfus Med Hemother.* 2013 Jun;40(3):172-81.