

## Del – das weak D Ostasiens?

1. Okubo Y, Yamaguchi H, Tomita T, Nagao N. A D variant, Del? *Transfusion*. 1984;24(6):542.
2. Garratty G. The significance of IgG on the red cell surface. *Transfus Med Rev*. 1987;1(1):47-57.
3. Fukumori Y, Hori Y, Ohnoki S, Nagao N, Shibata H, Okubo Y, et al. Further analysis of Del (D-elute) using polymerase chain reaction (PCR) with RHD gene-specific primers. *Transfus Med*. 1997;7(3):227-31.
4. Chang J-G, Wang J-C, Yang T-Y, Tsan K-W, Shih M-C, Peng C-T, et al. Human RhDel Is Caused by a Deletion of 1,013 bp Between Introns 8 and 9 Including Exon 9 of RHD Gene. *Blood*. 1998;92(7):2602-4.
5. Wagner F, Frohmajer A, Flegel WA. RHD positive haplotypes in D negative Europeans. *BMC Genetics*. 2001;2:15.
6. Yin Q, Flegel WA. DEL in China: the D antigen among serologic RhD-negative individuals. *J Transl Med*. 2021;19(1):439.
7. Choi Y, Sims GE, Murphy S, Miller JR, Chan AP. Predicting the functional effect of amino acid substitutions and indels. *PLoS One*. 2012;7(10):e46688.
8. Christiansen M, Sorensen BS, Grunnet N. RHD positive among C/E+ and D- blood donors in Denmark. *Transfusion*. 2010;50(7):1460-4.
9. Shao CP. Transfusion of RhD-positive blood in "Asia type" DEL recipients. *N Engl J Med*. 2010;362(5):472-3.
10. Wang QP, Dong GT, Wang XD, Gu J, Li Z, Sun AY, et al. An investigation of secondary anti-D immunisation among phenotypically RhD-negative individuals in the Chinese population. *Blood Transfus*. 2014;12(2):238-43.
11. Wang M, Wang BL, Xu W, Fan DD, Peng ML, Pan J, et al. Anti-D alloimmunisation in pregnant women with DEL phenotype in China. *Transfus Med*. 2015;25(3):163-9.
12. Xu W, Zhu M, Wang BL, Su H, Wang M. Prospective Evaluation of a Transfusion Policy of RhD-Positive Red Blood Cells into DEL Patients in China. *Transfus Med Hemother*. 2015;42(1):15-21.
13. Ito S, Ohto H, Ogiyama Y, Irino M, Omokawa S, Shibasaki I, et al. A practical and effective strategy in East Asia to prevent anti D alloimmunization in patients by C/c phenotyping of serologic RhD negative blood donors. *eJHaem*. 2021;2(4):750-6.
14. Gassner C, Doescher A, Drnovsek TD, Rozman P, Eicher NI, Legler TJ, et al. Presence of RHD in serologically D-, C/E+ individuals: a European multicenter study. *Transfusion*. 2005;45(4):527-38.
15. Krog GR, Clausen FB, Berkowicz A, Jorgensen L, Rieneck K, Nielsen LK, et al. Is current serologic RhD typing of blood donors sufficient for avoiding immunization of recipients? *Transfusion*. 2011;51(11):2278-85.
16. Shao CP, Wang BY, Ye SH, Zhang WL, Xu H, Zhuang NB, et al. DEL RBC transfusion should be avoided in particular blood recipient in East Asia due to allosensitization and ineffectiveness. *J Zhejiang Univ Sci B*. 2012;13(11):913-8.
17. St-Louis M, Lebrun A, Goldman M, Lavoie M. Alloimmunization of patients by blood units harboring distinct DEL variants. *Immunohematology*. 2013;29(4):136-40.
18. Perez-Alvarez I, Hayes C, Hailemariam T, Shin E, Hutchinson T, Klapper E. RHD genotyping of serologic RhD-negative blood donors in a hospital-based blood donor center. *Transfusion*. 2019;59(7):2422-8.
19. Gowland P, Gassner C, Hustinx H, Stolz M, Gottschalk J, Tissot JD, et al. Molecular RHD screening of RhD negative donors can replace standard serological testing for RhD negative donors. *Transfusion and apheresis science : official journal of the World Apheresis Association : official journal of the European Society for Haemapheresis*. 2014;50(2):163-8.
20. Lejon Crottet S, Haer-Wigman L, Gowland P, Fontana S, Niederhauser C, Hustinx H. Serologic and molecular investigations of DAR1 (weak D Type 4.2), DAR1.2, DAR1.3, DAR2 (DARE), and DARA. *Transfusion*. 2013;53(11 Suppl 2):3000-8.
21. Flegel WA, von Zabern I, Wagner FF. Six years' experience performing RHD genotyping to confirm D- red blood cell units in Germany for preventing anti-D immunizations. *Transfusion*. 2009;49(3):465-71.
22. Polin H, Danzer M, Gaszner W, Broda D, St-Louis M, Proll J, et al. Identification of RHD alleles with the potential of anti-D immunization among seemingly D- blood donors in Upper Austria. *Transfusion*. 2009;49(4):676-81.
23. Wagner FF, Frohmajer A, Ladewig B, Eicher NI, Lonicer CB, Muller TH, et al. Weak D alleles express distinct phenotypes. *Blood*. 2000;95(8):2699-708.