

Massivtransfusion – ein Update

1. Amgalan A, Allen T, Othman M et al. (2020) Systematic review of viscoelastic testing (TEG/ROTEM) in obstetrics and recommendations from the women's SSC of the ISTH. *J Thromb Haemost* 18:1813-1838. DOI: 10.1111/jth.14882.
2. Arbeitsgemeinschaft Der Wissenschaftlichen Medizinischen Fachgesellschaften e.V. (2023) AWMF-Leitlinie 187-023: überarbeitete S3-Leitlinie Polytrauma / Schwerverletzten-Behandlung. https://register.awmf.org/assets/guidelines/187-023_S3_Polytrauma-Schwerverletzten-Behandlung_2023-02.pdf. Zugegriffen: Feb 14, 2023.
3. Arbeitsgemeinschaft Der Wissenschaftlichen Medizinischen Fachgesellschaften e.V. (AWMF) (2022) AWMF-Leitlinie 015-063: Peripartale Blutungen, Diagnostik und Therapie. <https://www.awmf.org/leitlinien/detail/II/015-063.html>. Zugegriffen: Oct 29, 2022.
4. Badami KG, Neal C, Sparrow RL et al. (2022) Red blood cell alloantibodies in the context of critical bleeding and massive transfusion. *Blood Transfus* DOI: 10.2450/2022.0131-22.
5. Baksaas-Aasen K, Van Dieren S, Balvers K et al. (2019) Data-driven Development of ROTEM and TEG Algorithms for the Management of Trauma Hemorrhage: A Prospective Observational Multicenter Study. *Ann Surg* 270:1178-1185. DOI: 10.1097/SLA.0000000000002825.
6. Beekley AC (2008) Damage control resuscitation: a sensible approach to the exsanguinating surgical patient. *Crit Care Med* 36:S267-274. DOI: 10.1097/CCM.0b013e31817da7dc.
7. Bogner-Flatz V, Braunstein M, Ocker LE et al. (2019) On-the-Scene Hyaluronan and Syndecan-1 Serum Concentrations and Outcome after Cardiac Arrest and Resuscitation. *Mediators Inflamm* 2019:8071619. DOI: 10.1155/2019/8071619.
8. Bonanno FG (2022) Management of Hemorrhagic Shock: Physiology Approach, Timing and Strategies. *J Clin Med* 12 DOI: 10.3390/jcm12010260.
9. Bossers SM, Loer SA, Bloemers FW et al. (2021) Association Between Prehospital Tranexamic Acid Administration and Outcomes of Severe Traumatic Brain Injury. *JAMA Neurol* 78:338-345. DOI: 10.1001/jamaneurol.2020.4596.
10. Brohi K, Singh J, Heron M et al. (2003) Acute traumatic coagulopathy. *J Trauma* 54:1127-1130. DOI: 10.1097/01.TA.0000069184.82147.06.
11. Bundesärztekammer (BÄK, German Medical Association), (2017) Richtlinie zur Gewinnung von Blut und Blutbestandteilen und zur Anwendung von Blutprodukten (Richtlinie Hämotherapie) Gesamtnovelle 2017. <http://www.bundesaerztekammer.de/aerzte/medizin-ethik/wissenschaftlicher-beirat/veroeffentlichungen/haemotherapie-transfusionsmedizin/richtlinie>. Zugegriffen: Aug 06, 2017.
12. Bundesärztekammer (BÄK) (2020) Querschnitts-Leitlinien (BÄK) zur Therapie mit Blutkomponenten und Plasmaderivaten – Gesamtnovelle 2020. <https://www.bundesaerztekammer.de/aerzte/medizin-ethik/wissenschaftlicher-beirat/veroeffentlichungen/haemotherapietransfusionsmedizin/querschnitt-leitlinie>. Zugegriffen: Jan 10, 2021.
13. Chowdary P, Saayman AG, Paulus U et al. (2004) Efficacy of standard dose and 30 ml/kg fresh frozen plasma in correcting laboratory parameters of haemostasis in critically ill patients. *Br J Haematol* 125:69-73. DOI: 10.1111/j.1365-2141.2004.04868.x.
14. Coleman JR, Moore EE, Samuels JM et al. (2021) Whole Blood Thrombin Generation in Severely Injured Patients Requiring Massive Transfusion. *J Am Coll Surg* 232:709-716. DOI: 10.1016/j.jamcollsurg.2020.12.058.
15. Collins PW, Cannings-John R, Bruynseels D et al. (2017) Viscoelastometric-guided early fibrinogen concentrate replacement during postpartum haemorrhage: OBS2, a double-blind randomized controlled trial. *Br J Anaesth* 119:411-421. DOI: 10.1093/bja/aex181.
16. Cusack R, Leone M, Rodriguez AH et al. (2022) Endothelial Damage and the Microcirculation in Critical Illness. *Biomedicines* 10 DOI: 10.3390/biomedicines10123150.
17. Debot M, Eitel AP, Moore EE et al. (2022) Blood Type O Is a Risk Factor for Hyperfibrinolysis and Massive Transfusion after Severe Injury. *Shock* 58:492-497. DOI: 10.1097/SHK.0000000000002013.
18. Derickson MJ, McClellan JM, Marko ST et al. (2018) The effects of hemorrhage on the pharmacokinetics of tranexamic acid in a swine model. *J Trauma Acute Care Surg* 85:S44-S48. DOI: 10.1097/TA.0000000000001861.
19. Dettmer M, Morrison J, Bari V et al. (2022) Factors Associated With Mortality Among Patients Managed for Large Volume Hemorrhage in a Medical Intensive Care Unit. *Shock* 57:392-396. DOI: 10.1097/SHK.0000000000001913.
20. Ditzel RM, Jr., Anderson JL, Eisenhart WJ et al. (2020) A review of transfusion- and trauma-induced hypocalcemia: Is it time to change the lethal triad to the lethal diamond? *J Trauma Acute Care Surg* 88:434-439. DOI: 10.1097/TA.0000000000002570.
21. Dorken-Gallastegi A, Renne AM, Bokenkamp M et al. (2022) Balanced blood component resuscitation in trauma: Does it matter equally at different transfusion volumes? *Surgery* DOI: 10.1016/j.surg.2022.11.010.
22. Drukker L, Srebni N, Elstein D et al. (2016) The association between ABO blood group and obstetric hemorrhage. *J Thromb Thrombolysis* 42:340-345. DOI: 10.1007/s11239-016-1360-5.

23. Dunbar NM, Yazer MH, Biomedical Excellence for Safer Transfusion C et al. (2017) Safety of the use of group A plasma in trauma: the STAT study. *Transfusion* 57:1879-1884. DOI: 10.1111/trf.14139.
24. Dutton RP (2012) Haemostatic resuscitation. *Br J Anaesth* 109 Suppl 1:i39-i46. DOI: 10.1093/bja/aes389.
25. El-Menyar A, Ahmed K, Hakim S et al. (2022) Efficacy and safety of the second in-hospital dose of tranexamic acid after receiving the prehospital dose: double-blind randomized controlled clinical trial in a level 1 trauma center. *Eur J Trauma Emerg Surg* 48:3089-3099. DOI: 10.1007/s00068-021-01848-0.
26. Erdoes G, Koster A, Ortmann E et al. (2021) A European consensus statement on the use of four-factor prothrombin complex concentrate for cardiac and non-cardiac surgical patients. *Anaesthesia* 76:381-392. DOI: 10.1111/anae.15181.
27. Franchini M, Favaloro EJ, Targher G et al. (2012) ABO blood group, hypercoagulability, and cardiovascular and cancer risk. *Crit Rev Clin Lab Sci* 49:137-149. DOI: 10.3109/10408363.2012.708647.
28. Franco L, Becattini C, Beyer-Westendorf J et al. (2020) Definition of major bleeding: Prognostic classification. *J Thromb Haemost* 18:2852-2860. DOI: 10.1111/jth.15048.
29. Frietsch T, Steinbicker AU, Horn A et al. (2022) Safety of Intraoperative Cell Salvage in Cancer Surgery: An Updated Meta-Analysis of the Current Literature. *Transfusion Medicine and Hemotherapy* 49:143-157. DOI: 10.1159/000524538.
30. Funk M, Heiden M, Volz-Zang C (2017) Hämovigilanz-Bericht des Paul-Ehrlich-Institutes 2015. Auswertung der Meldungen von schweren Reaktionen und Zwischenfällen nach §63i AMG. <http://www.pei.de/DE/arzneimittelsicherheit-vigilanz/haemovigilanz/haemovigilanzberichte/haemovigilanzberichte-node.html>. Zugegriffen: Jul 02, 2017.
31. Gelbard RB, Griffin RL, Reynolds L et al. (2022) Over-transfusion with blood for suspected hemorrhagic shock is not associated with worse clinical outcomes. *Transfusion* 62 Suppl 1:S177-S184. DOI: 10.1111/trf.16978.
32. Gernsheimer TB, Brown SP, Triulzi DJ et al. (2022) Prophylactic tranexamic acid in patients with hematologic malignancy: a placebo-controlled, randomized clinical trial. *Blood* 140:1254-1262. DOI: 10.1182/blood.2022016308.
33. Gianola S, Castellini G, Biffi A et al. (2022) Accuracy of risk tools to predict critical bleeding in major trauma: A systematic review with meta-analysis. *J Trauma Acute Care Surg* 92:1086-1096. DOI: 10.1097/TA.0000000000003496.
34. Gill JC, Endres-Brooks J, Bauer PJ et al. (1987) The effect of ABO blood group on the diagnosis of von Willebrand disease. *Blood* 69:1691-1695.
35. Guerado E, Medina A, Mata MI et al. (2016) Protocols for massive blood transfusion: when and why, and potential complications. *Eur J Trauma Emerg Surg* 42:283-295. DOI: 10.1007/s00068-015-0612-y.
36. Gustafson ML, Hollosi S, Chumbe JT et al. (2015) The effect of ethanol on lactate and base deficit as predictors of morbidity and mortality in trauma. *Am J Emerg Med* 33:607-613. DOI: 10.1016/j.ajem.2015.01.030.
37. Halt-It Trial Collaborators (2020) Effects of a high-dose 24-h infusion of tranexamic acid on death and thromboembolic events in patients with acute gastrointestinal bleeding (HALT-IT): an international randomised, double-blind, placebo-controlled trial. *Lancet* 395:1927-1936. DOI: 10.1016/S0140-6736(20)30848-5.
38. Hamada SR, Garrigue D, Nougue H et al. (2022) Impact of platelet transfusion on outcomes in trauma patients. *Crit Care* 26:49. DOI: 10.1186/s13054-022-03928-y.
39. He Q, Zhou Y, Wang F et al. (2019) Blood Type O Predicts Hematoma Expansion in Patients with Intracerebral Hemorrhage. *J Stroke Cerebrovasc Dis* 28:2311-2317. DOI: 10.1016/j.jstrokecerebrovasdis.2019.05.022.
40. Hess JR, Holcomb JB (2015) Resuscitating PROPPRly. *Transfusion* 55:1362-1364. DOI: 10.1111/trf.13118.
41. Holcomb JB, Tilley BC, Baraniuk S et al. (2015) Transfusion of plasma, platelets, and red blood cells in a 1:1:1 vs a 1:1:2 ratio and mortality in patients with severe trauma: the PROPPR randomized clinical trial. *JAMA* 313:471-482. DOI: 10.1001/jama.2015.12.
42. Honickel M, Treutler S, Van Ryn J et al. (2015) Reversal of dabigatran anticoagulation ex vivo: Porcine study comparing prothrombin complex concentrates and idarucizumab. *Thromb Haemost* 113:728-740. DOI: 10.1160/TH14-08-0712.
43. Innerhofer N, Treichl B, Rugg C et al. (2021) First-Line Administration of Fibrinogen Concentrate in the Bleeding Trauma Patient: Searching for Effective Dosages and Optimal Post-Treatment Levels Limiting Massive Transfusion-Further Results of the RETIC Study. *J Clin Med* 10DOI: 10.3390/jcm10173930.
44. Innerhofer P, Fries D, Mittermayr M et al. (2017) Reversal of trauma-induced coagulopathy using first-line coagulation factor concentrates or fresh frozen plasma (RETIC): a single-centre, parallel-group, open-label, randomised trial. *The Lancet Haematology* 4:e258-e271. DOI: 10.1016/s2352-3026(17)30077-7.
45. Johansson PI (2010) Goal-directed hemostatic resuscitation for massively bleeding patients: the Copenhagen concept. *Transfus Apher Sci* 43:401-405. DOI: 10.1016/j.transci.2010.09.002.

46. Joint United Kingdom (UK) (2020) Blood Transfusion and Tissue Transplantation Services Professional Advisory Committee (Jpac) 7.3: Transfusion management of major haemorrhage. <https://www.transfusionguidelines.org/transfusion-handbook/7-effective-transfusion-in-surgery-and-critical-care/7-3-transfusion-management-of-major-haemorrhage>. Zugegriffen: Jan 20, 2023.
47. Kahr MK, Franke D, Brun R et al. (2018) Blood group O: A novel risk factor for increased postpartum blood loss? *Haemophilia* 24:e207-e212. DOI: 10.1111/hae.13537.
48. Kalbhenn J, Wittau N, Schmutz A et al. (2015) Identification of acquired coagulation disorders and effects of target-controlled coagulation factor substitution on the incidence and severity of spontaneous intracranial bleeding during veno-venous ECMO therapy. *Perfusion* 30:675-682. DOI: 10.1177/0267659115579714.
49. Kamyszek RW, Leraas HJ, Reed C et al. (2019) Massive transfusion in the pediatric population: A systematic review and summary of best-evidence practice strategies. *J Trauma Acute Care Surg* 86:744-754. DOI: 10.1097/TA.0000000000002188.
50. Kao TW, Lee YC, Chang HT (2021) Prothrombin Complex Concentrate for Trauma Induced Coagulopathy: A Systematic Review and Meta-Analysis. *J Acute Med* 11:81-89. DOI: 10.6705/j.jacme.202109_11(3).0001.
51. Kashuk JL, Moore EE, Millikan JS et al. (1982) Major abdominal vascular trauma--a unified approach. *J Trauma* 22:672-679. DOI: 10.1097/00005373-198208000-00004.
52. Klein AA, Arnold P, Bingham RM et al. (2016) AAGBI guidelines: the use of blood components and their alternatives 2016. *Anaesthesia* 71:829-842. DOI: 10.1111/anae.13489.
53. Kogutt BK, Vaught AJ (2019) Postpartum hemorrhage: Blood product management and massive transfusion. *Seminars in perinatology* 43:44-50. DOI: 10.1053/j.semperi.2018.11.008.
54. Kolev K, Longstaff C (2016) Bleeding related to disturbed fibrinolysis. *Br J Haematol* 175:12-23. DOI: 10.1111/bjh.14255.
55. Koscielny J, Von Tempelhoff GF, Ziemer S et al. (2004) A practical concept for preoperative management of patients with impaired primary hemostasis. *Clin Appl Thromb Hemost* 10:155-166.
56. Kozek-Langenecker SA, Ahmed AB, Afshari A et al. (2017) Management of severe perioperative bleeding: guidelines from the European Society of Anaesthesiology: First update 2016. *Eur J Anaesthesiol* 34:332-395. DOI: 10.1097/EJA.0000000000000630.
57. Kregel HR, Hatton GE, Isbell KD et al. (2022) Shock-Induced Endothelial Dysfunction is Present in Patients With Occult Hypoperfusion After Trauma. *Shock* 57:106-112. DOI: 10.1097/SHK.0000000000001866.
58. Lawati KA, Sharif S, Maqbali SA et al. (2021) Efficacy and safety of tranexamic acid in acute traumatic brain injury: a systematic review and meta-analysis of randomized-controlled trials. *Intensive Care Med* 47:14-27. DOI: 10.1007/s00134-020-06279-w.
59. Lier H, Gathof B (2017) Massivtransfusion – logistische und therapeutische Aspekte. *Hämotherapie* 29/2017:1-5.
60. Lier H, Michael M (2019) Notaufnahme: Management starker Blutungen. *Notaufnahme up2date* 1:47-63. DOI: 10.1055/a-0926-0978.
61. Lv K, Yuan Q, Fu P et al. (2020) Impact of fibrinogen level on the prognosis of patients with traumatic brain injury: a single-center analysis of 2570 patients. *World J Emerg Surg* 15:54. DOI: 10.1186/s13017-020-00332-1.
62. McLaughlin DF, Niles SE, Salinas J et al. (2008) A predictive model for massive transfusion in combat casualty patients. *J Trauma* 64:S57-63; discussion S63. DOI: 10.1097/TA.0b013e318160a566.
63. Meretoja A, Yassi N, Wu TY et al. (2020) Tranexamic acid in patients with intracerebral haemorrhage (STOP-AUST): a multicentre, randomised, placebo-controlled, phase 2 trial. *Lancet Neurol* 19:980-987. DOI: 10.1016/S1474-4422(20)30369-0.
64. Meyer DE, Cotton BA, Fox EE et al. (2018) A comparison of resuscitation intensity and critical administration threshold in predicting early mortality among bleeding patients: A multicenter validation in 680 major transfusion patients. *J Trauma Acute Care Surg* 85:691-696. DOI: 10.1097/TA.0000000000002020.
65. Mikhail J (1999) The trauma triad of death: hypothermia, acidosis, and coagulopathy. *AACN Clin Issues* 10:85-94.
66. Miller RD, Robbins TO, Tong MJ et al. (1971) Coagulation defects associated with massive blood transfusions. *Ann Surg* 174:794-801. DOI: 10.1097/0000658-197111000-00010.
67. Moore EE (1996) Thomas G. Orr Memorial Lecture. Staged laparotomy for the hypothermia, acidosis, and coagulopathy syndrome. *Am J Surg* 172:405-410. DOI: 10.1016/s0002-9610(96)00216-4.
68. Moore SA, Raval JS (2022) Massive transfusion: a review. *Ann Blood* 7:18. DOI: 10.21037/aob-22-3.
69. Nascimento B, Callum J, Tien H et al. (2016) Fibrinogen in the initial resuscitation of severe trauma (FiiRST): a randomized feasibility trial. *Br J Anaesth* 117:775-782. DOI: 10.1093/bja/aew343.

70. Naumann DN, Hazeldine J, Davies DJ et al. (2018) Endotheliopathy of Trauma is an on-Scene Phenomenon, and is Associated with Multiple Organ Dysfunction Syndrome: A Prospective Observational Study. *Shock* 49:420-428. DOI: 10.1097/shk.0000000000000999.
71. Neeki MM, Dong F, Toy J et al. (2017) Efficacy and Safety of Tranexamic Acid in Prehospital Traumatic Hemorrhagic Shock: Outcomes of the Cal-PAT Study. *West J Emerg Med* 18:673-683. DOI: 10.5811/westjem.2017.2.32044.
72. Nunez TC, Voskresensky IV, Dossett LA et al. (2009) Early prediction of massive transfusion in trauma: simple as ABC (assessment of blood consumption)? *J Trauma* 66:346-352. DOI: 10.1097/TA.0b013e3181961c35.
73. Owattanapanich N, Chittawatanarat K, Benyakorn T et al. (2018) Risks and benefits of hypotensive resuscitation in patients with traumatic hemorrhagic shock: a meta-analysis. *Scand J Trauma Resusc Emerg Med* 26:107. DOI: 10.1186/s13049-018-0572-4.
74. Pommerening MJ, Goodman MD, Holcomb JB et al. (2015) Clinical gestalt and the prediction of massive transfusion after trauma. *Injury* 46:807-813. DOI: 10.1016/j.injury.2014.12.026.
75. Ponschab M, Schochl H, Gabriel C et al. (2015) Haemostatic profile of reconstituted blood in a proposed 1:1:1 ratio of packed red blood cells, platelet concentrate and four different plasma preparations. *Anaesthesia* 70:528-536. DOI: 10.1111/anae.13067.
76. Post R, Germans MR, Tjerkstra MA et al. (2021) Ultra-early tranexamic acid after subarachnoid haemorrhage (ULTRA): a randomised controlled trial. *Lancet* 397:112-118. DOI: 10.1016/S0140-6736(20)32518-6.
77. Qin X, Zhang W, Zhu X et al. (2021) Early Fresh Frozen Plasma Transfusion: Is It Associated With Improved Outcomes of Patients With Sepsis? *Front Med (Lausanne)* 8:754859. DOI: 10.3389/fmed.2021.754859.
78. Quintana MT, Zebley JA, Vincent A et al. (2022) Cresting mortality: Defining a plateau in ongoing massive transfusion. *J Trauma Acute Care Surg* 93:43-51. DOI: 10.1097/TA.0000000000003641.
79. Rahbar E, Fox EE, Del Junco DJ et al. (2013) Early resuscitation intensity as a surrogate for bleeding severity and early mortality in the PROMMTT study. *J Trauma Acute Care Surg* 75:S16-23. DOI: 10.1097/TA.0b013e31828fa535.
80. Richards JE, Harris T, Dunser MW et al. (2021) Vasopressors in Trauma: A Never Event? *Anesth Analg* 133:68-79. DOI: 10.1213/ANE.0000000000005552.
81. Rizoli S, Min A, Sanchez AP et al. (2016) In Trauma, Conventional ROTEM and TEG Results Are Not Interchangeable But Are Similar in Clinical Applicability. *Mil Med* 181:117-126. DOI: 10.7205/MILMED-D-15-00166.
82. Rosenfeld E, Lau P, Zhang W et al. (2019) Defining massive transfusion in civilian pediatric trauma. *J Pediatr Surg* 54:975-979. DOI: 10.1016/j.jpedsurg.2019.01.029.
83. Sabouri M, Vahidian M, Sourani A et al. (2022) Efficacy and safety of fibrinogen administration in acute post-traumatic hypofibrinogenemia in isolated severe traumatic brain injury: A randomized clinical trial. *J Clin Neurosci* 101:204-211. DOI: 10.1016/j.jocn.2022.05.016.
84. Sagraves SG, Toschlog EA, Rotondo MF (2006) Damage control surgery--the intensivist's role. *J Intensive Care Med* 21:5-16. DOI: 10.1177/0885066605282790.
85. Sanghavi M, Rutherford JD (2014) Cardiovascular physiology of pregnancy. *Circulation* 130:1003-1008. DOI: 10.1161/CIRCULATIONAHA.114.009029.
86. Savage SA, Sumislawski JJ, Zarzaur BL et al. (2015) The new metric to define large-volume hemorrhage: results of a prospective study of the critical administration threshold. *J Trauma Acute Care Surg* 78:224-229; discussion 229-230. DOI: 10.1097/TA.0000000000000502.
87. Savage SA, Zarzaur BL, Croce MA et al. (2013) Redefining massive transfusion when every second counts. *J Trauma Acute Care Surg* 74:396-400; discussion 400-392. DOI: 10.1097/TA.0b013e31827a3639.
88. Sawamura A, Hayakawa M, Gando S et al. (2009) Disseminated intravascular coagulation with a fibrinolytic phenotype at an early phase of trauma predicts mortality. *Thromb Res* 124:608-613. DOI: 10.1016/j.thromres.2009.06.034.
89. Schlimp CJ, Schochl H (2014) The role of fibrinogen in trauma-induced coagulopathy. *Hamostaseologie* 34:29-39. DOI: 10.5482/HAMO-13-07-0038.
90. Schochl H, Voelckel W, Maegele M et al. (2014) Endogenous thrombin potential following hemostatic therapy with 4-factor prothrombin complex concentrate: a 7-day observational study of trauma patients. *Crit Care* 18:R147. DOI: 10.1186/cc13982.
91. Schroll R, Swift D, Tatum D et al. (2018) Accuracy of shock index versus ABC score to predict need for massive transfusion in trauma patients. *Injury* 49:15-19. DOI: 10.1016/j.injury.2017.09.015.
92. Schulman S, Kearon C, Subcommittee on Control of Anticoagulation of The S et al. (2005) Definition of major bleeding in clinical investigations of antihemostatic medicinal products in non-surgical patients. *J Thromb Haemost* 3:692-694. DOI: 10.1111/j.1538-7836.2005.01204.x.
93. Seheult JN, Dunbar NM, Hess JR et al. (2020) Transfusion of blood components containing ABO-incompatible plasma does not lead to higher mortality in civilian trauma patients. *Transfusion* 60:2517-2528. DOI: 10.1111/trf.16008.

94. Shah A, Kerner V, Stanworth SJ et al. (2023) Major haemorrhage: past, present and future. *Anaesthesia* 78:93-104. DOI: 10.1111/anae.15866.
95. Spahn DR, Bouillon B, Cerny V et al. (2019) The European guideline on management of major bleeding and coagulopathy following trauma: fifth edition. *Crit Care* 23:98. DOI: 10.1186/s13054-019-2347-3.
96. Sprigg N, Flaherty K, Appleton JP et al. (2018) Tranexamic acid for hyperacute primary IntraCerebral Haemorrhage (TICH-2): an international randomised, placebo-controlled, phase 3 superiority trial. *Lancet* 391:2107-2115. DOI: 10.1016/S0140-6736(18)31033-X.
97. Stanworth SJ, Dowling K, Curry N et al. (2022) Haematological management of major haemorrhage: a British Society for Haematology Guideline. *Br J Haematol* 198:654-667. DOI: 10.1111/bjh.18275.
98. Tachon G, Harrois A, Tanaka S et al. (2014) Microcirculatory alterations in traumatic hemorrhagic shock. *Crit Care Med* 42:1433-1441. DOI: 10.1097/CCM.000000000000223.
99. Tieu BH, Holcomb JB, Schreiber MA (2007) Coagulopathy: its pathophysiology and treatment in the injured patient. *World J Surg* 31:1055-1064. DOI: 10.1007/s00268-006-0653-9.
100. Tran A, Yates J, Lau A et al. (2018) Permissive hypotension versus conventional resuscitation strategies in adult trauma patients with hemorrhagic shock: A systematic review and meta-analysis of randomized controlled trials. *J Trauma Acute Care Surg* 84:802-808. DOI: 10.1097/TA.0000000000001816.
101. Van Den Brink DP, Wirtz MR, Neto AS et al. (2020) Effectiveness of prothrombin complex concentrate for the treatment of bleeding: A systematic review and meta-analysis. *J Thromb Haemost* 18:2457-2467. DOI: 10.1111/jth.14991.
102. Van Wessel KJP, Jochems D, Leenen LPH (2022) The effect of prehospital tranexamic acid on outcome in polytrauma patients with associated severe brain injury. *Eur J Trauma Emerg Surg* 48:1589-1599. DOI: 10.1007/s00068-021-01827-5.
103. Vang M, Ostberg M, Steinmetz J et al. (2022) Shock index as a predictor for mortality in trauma patients: a systematic review and meta-analysis. *Eur J Trauma Emerg Surg* 48:2559-2566. DOI: 10.1007/s00068-022-01932-z.
104. Watson GA, Sperry JL, Rosengart MR et al. (2009) Fresh frozen plasma is independently associated with a higher risk of multiple organ failure and acute respiratory distress syndrome. *J Trauma* 67:221-227; discussion 228-230. DOI: 10.1097/TA.0b013e3181ad5957.
105. Wu F, Kozar RA (2019) Fibrinogen Protects Against Barrier Dysfunction Through Maintaining Cell Surface Syndecan-1 In Vitro. *Shock* 51:740-744. DOI: 10.1097/SHK.0000000000001207.
106. Yucel N, Lefering R, Maegele M et al. (2006) Trauma Associated Severe Hemorrhage (TASH)-Score: probability of mass transfusion as surrogate for life threatening hemorrhage after multiple trauma. *J Trauma* 60:1228-1236. DOI: 10.1097/01.ta.0000220386.84012.bf.
107. Zhu CS, Braverman M, Goddard S et al. (2021) Prehospital shock index and systolic blood pressure are highly specific for pediatric massive transfusion. *J Trauma Acute Care Surg* 91:579-583. DOI: 10.1097/TA.0000000000003275.
108. Zipperle J, Ziegler B, Schochl H et al. (2022) Conventional and Pro-Inflammatory Pathways of Fibrinolytic Activation in Non-Traumatic Hyperfibrinolysis. *J Clin Med* 11. DOI: 10.3390/jcm11247305.