

Der Hämoglobinwert als Transfusionstrigger für Erythrozytenkonzentrate: Review und Nutzenbewertung randomisierter klinischer Studien

1. Hebert PC, Wells G, Blajchman MA, Marshall J, Martin, C, Pagliarello G, Tweeddale M, Schweitzer I, Yetisir E. A multicenter, randomized, controlled clinical trial of transfusion requirements in critical care. *The New England Journal of Medicine* 1999;340: 409-17.
2. Carson JL, Duff A, Poses RM, Berlin JA, Spence RK, Trout R, Noveck H, Strom BL. Effect of anaemia and cardiovascular disease on surgical mortality and morbidity. *Lancet* 1996;348:1055-60.
3. Hebert PC, Wells G, Tweeddale M, et al. Does transfusion practice affect mortality in critically ill patients? *Am J Respir Crit Care Med* 1997;155:1618-23.
4. Carson JL, Brooks MM, Abbott JD, Chaitman B, Kelsey SF, Triulzi DJ, Srinivas V, Menegus MA, Marroquin OC, Rao SV, Noveck H, Passano E, Hardison RM, Smitherman T, Vagaonescu T, Wimmer NJ, Williams DO. Liberal versus restrictive transfusion thresholds for patients with symptomatic coronary artery disease. *American heart journal* 2013;165: 964-71 e1.
5. Lacroix J, Hebert PC, Hutchison JS, Hume HA, Tucci M, Ducruet T, Gauvin F, Collet JP, Toledano BJ, Robillard P, Joffe A, Biarent D, Meert K, Peters MJ. Transfusion strategies for patients in pediatric intensive care units. *The New England Journal of Medicine* 2007;356:1609-19.
6. Holst LB, Haase N, Wetterslev J, Wernerman J, Guttormsen AB, et al. Lower versus higher hemoglobin threshold for transfusion in septic shock. *The New England Journal of Medicine* 2014;371:1381-91.
7. Villanueva C, Colomo A, Bosch A, Concepcion M, Hernandez-Gea V, Aracil C, Graupera I, Poca M, Alvarez-Urturi C, Gordillo J, Guarner-Argente C, Santalo M, Muniz E, Guarner C. Transfusion strategies for acute upper gastrointestinal bleeding. *The New England journal of medicine* 2013;368: 11-21.
8. Hajjar LA, Vincent JL, Galas FRBG, Nakamura RE, Silva CMP, Santos MH, Fukushima J, et al. Transfusion requirements after cardiac surgery. *JAMA* 2010;304:1559-67.
9. Murphy GJ, Pike K, Rogers CA, Wordsworth S, Stokes EA, Angelini GD, Reeves BC. Liberal or restrictive transfusion after cardiac surgery. *The New England Journal of Medicine* 2015;372:997-1008.
10. Carson JL, Terrin ML, Noveck H, Sanders DW, Chaitman BR, Rhoads GG, Nemo G, Dragert K, Beaupre L, Hildebrand K, Macaulay W, Lewis C, Cook DR, Dobbin G, Zakriya KJ, Apple FS, Horney RA, Magaziner J. Liberal or restrictive transfusion in high-risk patients after hip surgery. *The New England journal of medicine* 2011;365: 2453-62.
11. Carson JL, Sieber F, Cook DR, Hoover DR, Noveck H, Chaitman BR, Fleisher L, Beaupre L, Macaulay W, Rhoads GG, Paris B, Zagorin A, Sanders DW, Zakriya KJ, Magaziner J. Liberal versus restrictive blood transfusion strategy: 3-year survival and cause of death results from the FOCUS randomised controlled trial. *Lancet* 2015;385:1183-9.
12. Middelburg RA, van de Watering LM, van der Bom JG. Blood transfusions: good or bad? Confounding by indication, an underestimated problem in clinical transfusion research. *Transfusion* 2010;50: 1181-3.
13. Ferraris VA, Davenport DL, Saha SP, Austin PC, Zwischenberger JB. Surgical outcomes and transfusion of minimal amounts of blood in the operating room. *Archives of surgery* 2012;147: 49-55.
14. Yazer MH, Triulzi DJ. Things aren't always as they seem: what the randomized trials of red blood cell transfusion tell us about adverse outcomes (CME). *Transfusion* 2014;54: 3243-6.