

Klinische Studien zur Untersuchung von Nebenwirkungen länger gelagerter Erythrozytenkonzentrate

- 1) Martin CM, Sibbald WJ, Lu X, Hebert P, Schweitzer I. Age of transfused red blood cells is associated with ICU length of stay. *Clin Invest Med* 1994;**17**(Suppl 4):124.
- 2) Dessertaine G, Hammer L, Chenais F, Remy J, Schwebel C, Tabah A, ra-Somohano C, Bonadona A, Hamidfar-Roy R, Barnoud D, et al. [Does red blood cell storage time still influence ICU survival?]. *Transfus.Clin.Biol.* 2008 Sep;**15**(4):154-9.
- 3) Koch CG, Li L, Sessler DI, Figueroa P, Hoeltge GA, Mihajevic T, Blackstone EH. Duration of red-cell storage and complications after cardiac surgery. *N.Engl.J.Med.* 2008 Mar 20;**358**(12):1229-39.
- 4) Leal-Noval SR, Munoz-Gomez M, Arellano-Orden V, Marin-Caballos A, Amaya-Villar R, Marin A, Puppo-Moreno A, Ferrandiz-Millon C, Flores-Cordero JM, Murillo-Cabezas F. Impact of age of transfused blood on cerebral oxygenation in male patients with severe traumatic brain injury. *Crit Care Med.* 2008 Apr;**36**(4):1290-6.
- 5) Weinberg JA, McGwin G, Jr., Griffin RL, Huynh VQ, Cherry SA, III, Marques MB, Reiff DA, Kerby JD, Rue LW, III. Age of transfused blood: an independent predictor of mortality despite universal leukoreduction. *J.Trauma* 2008 Aug;**65**(2):279-82.
- 6) Weinberg JA, McGwin G, Jr., Marques MB, Cherry SA, III, Reiff DA, Kerby JD, Rue LW, III. Transfusions in the less severely injured: does age of transfused blood affect outcomes? *J.Trauma* 2008 Oct;**65**(4):794-8.
- 7) Yap CH, Lau L, Krishnaswamy M, Gaskell M, Yip M. Age of transfused red cells and early outcomes after cardiac surgery. *Ann.Thorac.Surg.* 2008 Aug;**86**(2):554-9.
- 8) Kneyber MC, Gazendam RP, Markhorst DG, Plotz FB. Length of storage of red blood cells does not affect outcome in critically ill children. *Intensive Care Med.* 2009 Jan;**35**(1):179-80.
- 9) Mynster T, Nielsen HJ. The impact of storage time of transfused blood on postoperative infectious complications in rectal cancer surgery. Danish RANX05 Colorectal Cancer Study Group. *Scand.J.Gastroenterol.* 2000;**35**(2):212-7.
- 10) Vandromme MJ, McGwin G, Jr., Marques MB, Kerby JD, Rue LW, III, Weinberg JA. Transfusion and pneumonia in the trauma intensive care unit: an examination of the temporal relationship. *J.Trauma* 2009 Jul;**67**(1):97-101.
- 11) van de Watering L, Lorinser J, Versteegh M, Westendorp R, Brand A. Effects of storage time of red blood cell transfusions on the prognosis of coronary artery bypass graft patients. *Transfusion* 2006 Oct;**46**(10):1712-8.
- 12) Leal-Noval SR, Jara-Lopez I, Garcia-Garmendia JL, Marin-Niebla A, Herruzo-Aviles A, Camacho-Larana P, Loscertales J. Influence of erythrocyte concentrate storage time on postsurgical morbidity in cardiac surgery patients. *Anesthesiology* 2003 Apr;**98**(4):815-22.
- 13) Offner PJ, Moore EE, Biffl WL, Johnson JL, Silliman CC. Increased rate of infection associated with transfusion of old blood after severe injury. *Arch.Surg.* 2002;**137**(6):711-6; discussion 716-7.
- 14) Zallen G, Offner PJ, Moore EE, Blackwell J, Ciesla DJ, Gabriel J, Denny C, Silliman CC. Age of transfused blood is an independent risk factor for postinjury multiple organ failure. *Am.J.Surg.* 1999;**178**(6):570-2.
- 15) Purdy FR, Tweeddale MG, Merrick PM. Association of mortality with age of blood transfused in septic ICU patients. *Can.J.Anaesth.* 1997;**44**(12):1256-61.
- 16) Spinella PC, Carroll CL, Staff I, Gross R, Mc QJ, Keibel L, Wade CE, Holcomb JB. Duration of red blood cell storage is associated with increased incidence of deep vein thrombosis and in hospital mortality in patients with traumatic injuries. *Crit Care* 2009;**13**(5):R151.